UPLIZNA®
inebilizumab-cdon

AFTER INFUSION CONSIDERATIONS

Here's a list of considerations to keep in mind after your infusion:

Watch out for any possible side effects. If you think you're experiencing one, tell your healthcare provider right away. You may report side effects to FDA at 1-800-FDA-1088.



Mark your next infusion in your calendar.

Write down anything you would change or bring to your next infusion.

Let your Patient Access Liaison (PAL) know how your infusion went*

Do something nice for yourself! Reward yourself for taking care of your health with a yummy meal, a good book, or any other treat that you enjoy.

*If you have an UPLIZNA prescription but haven't connected with a Patient Access Liaison (PAL) yet, call 1-833-842-8477 Monday-Friday, 8 AM-8 PM ET. The PAL is a dedicated support partner who helps investigate, explain, and educate on the steps in your treatment experience. Your PAL partners to support and champion you while accomplishing your treatment goals.

What is UPLIZNA® (inebilizumab-cdon)?

UPLIZNA is a prescription medicine used to treat adults with: Neuromyelitis optica spectrum disorder (NMOSD who are anti-aquaporin-4 (AQP4) antibody positive; Immunoglobulin G4-related disease (IgG4-RD); Generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) or anti-muscle specific tyrosine kinase (MuSK) antibody positive. It is not known if UPLIZNA is safe or effective in children.

IMPORTANT SAFETY INFORMATION

Who should not receive UPLIZNA?

You should not receive UPLIZNA if you have:

- had a life-threatening infusion reaction to UPLIZNA.
- an active hepatitis B virus infection.
- active or untreated inactive (latent) tuberculosis.

Please see additional Important Safety information on next page.

IMPORTANT SAFETY INFORMATION (CONT'D)

What is the most important information I should know about UPLIZNA?

UPLIZNA may cause serious side effects, including:

Infusion reactions. UPLIZNA can cause infusion reactions, including anaphylaxis, that can be serious or may cause you to be hospitalized. You will be monitored during your infusion and for at least 1 hour after each infusion of UPLIZNA for signs and symptoms of an infusion reaction. Tell your healthcare provider right away if you get any of these symptoms:

- headache
- nausea
- sleepiness
- shortness of breath
- fever
- muscle aches
- rash
- palpitations

If you develop an infusion reaction, your healthcare provider may need to stop or slow down the rate of your infusion and treat your symptoms.

Infections. Infections can happen during treatment with UPLIZNA. Tell your healthcare provider right away if you have an infection or get any of these symptoms:

- painful and frequent urination
- nasal congestion, runny nose, sore throat, fever, chills, cough, body aches

- UPLIZNA taken before or after other medicines that weaken the immune system may increase your risk of getting infections.
- **Hepatitis B virus (HBV) reactivation.** Before starting treatment with UPLIZNA, your healthcare provider will do blood tests to check for hepatitis B viral infection. If you have ever had hepatitis B virus infection, the hepatitis B virus may become active again during or after treatment with UPLIZNA. Hepatitis B virus becoming active again (called reactivation) may cause serious liver problems, including liver failure or death. Your healthcare provider will monitor you if you are at risk for hepatitis B virus reactivation during treatment and after you stop receiving UPLIZNA.
- **Progressive Multifocal Leukoencephalopathy (PML).** PML may happen with UPLIZNA. PML is a rare brain infection that leads to death or severe disability. Symptoms of PML may get worse over days to weeks. Call your healthcare provider right away if you get any of these symptoms:
  - weakness on one side of the body
  - loss of coordination in your arms and legs
  - changes in your vision
  - changes in thinking or memory
  - confusion
  - changes in your personality
- **Tuberculosis (TB).** TB is caused by an infection in the lungs. Before starting treatment with UPLIZNA, your healthcare provider will check to see if you are at risk for getting TB or have ever had TB.
- **Vaccinations.** Certain vaccines, called "live" or "live attenuated" vaccines, are not recommended in people receiving UPLIZNA. Talk to your healthcare provider before receiving any vaccinations. If you have a baby and you were receiving UPLIZNA during pregnancy, it is important to tell your baby's healthcare provider about your UPLIZNA use so they can decide when your baby should receive any vaccine.

Before receiving UPLIZNA, tell your healthcare provider about all of your medical conditions, including if you:

- have or think you have an infection.
- have ever taken, currently take, or plan to take medicines that affect your immune system, or other treatments for NMOSD, IgG4-RD, or gMG. These medicines may increase your risk of getting an infection.

Please see additional Important Safety information on next page.

UPLIZNA™
inebilizumab-cdon

IMPORTANT SAFETY INFORMATION (CONT'D)

Before receiving UPLIZNA, tell your healthcare provider about all of your medical conditions, including if you: (Cont'd)

- have or have ever had hepatitis B or are a carrier of the hepatitis B virus.
- have or have ever had tuberculosis.
- have had a recent vaccination or are scheduled to receive any vaccinations. You should receive any required vaccines at least 4 weeks before you start treatment with UPLIZNA.
- are pregnant or plan to become pregnant. It is not known if UPLIZNA will harm your unborn baby. Females should use birth control (contraception) during treatment with UPLIZNA and for 6 months after your last infusion of UPLIZNA.
- are breastfeeding or plan to breastfeed. It is not known if UPLIZNA passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you receive UPLIZNA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of UPLIZNA?

UPLIZNA may cause serious side effects, including:

- **low blood cell counts.** UPLIZNA may cause a decrease in some types of blood cells. Your healthcare provider will do blood tests to check your blood cell counts.

The most common side effects include: urinary tract infection and joint pain in patients with NMOSD; urinary tract infection and low white blood cell count in patients with IgG4-RD; headache and infusion-related reaction in patients with gMG. These are not all the possible side effects of UPLIZNA.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see UPLIZNA full Prescribing Information, including Medication Guide.

AMGEN © 2025 Amgen Inc. All rights reserved. USA-335-81489 12/25

UPLIZNA® inebilizumab-cdon